



1. PURPOSE

This Standard Operating Procedure (SOP) describes methods for anesthetizing rabbits.

2. RESPONSIBILITY

Principal investigator (PI) and their research staff.

3. GENERAL CONSIDERATIONS

- 3.1. Perform a thorough physical exam and obtain an accurate weight.
 - 3.2. Heat loss is rapid in anesthetized animals. Keep animals warm by providing a heat source until the animal has recovered from anesthesia.
 - 3.3. **Never leave an anesthetized animal unattended.** Monitor anesthetized animals until they fully recover, are sternal, and move into the cage.
-

4. MATERIALS

- 4.1. Material or equipment to provide or conserve body heat (e.g., heating disc or pad, warm-water circulating pad)
- 4.2. Ophthalmic ointment (natural tears) or surgical tape
- 4.3. Gas anesthesia machine (calibrated within the last 12 months) with adequate gas scavenging system or filter
- 4.4. Anesthesia mask
- 4.5. 2.5-3" endotracheal tube
- 4.6. Stethoscope with a hole in a plastic tube
- 4.7. Intra- vein catheter
- 4.8. Isoflurane
- 4.9. Ketamine (100mg/mL) *Controlled Drug
- 4.10. Xylazine (20mg/mL)
- 4.11. Acepromazine (10mg/mL)
- 4.12. Propofol (10mg/ml)
- 4.13. Sterile lubricant (e.g., water-soluble gel), 2% lidocaine gel

5. PROCEDURE

- 5.1. Apply ophthalmic ointment (natural tears) or surgical tape to both eyes to prevent dryness and damage to the cornea.
- 5.2. Inject Acepromazine 2mg/kg IM for tranquilization
- 5.3. Place an intravenous catheter (marginal ear vein):
 - 5.3.1. To provide IV fluid therapy and venous access during anesthesia.
 - 5.3.2. It is recommended to apply EMLA cream over the venipuncture site at least 15 minutes before placing the catheter.
- 5.4. Sedation:
 - 5.4.1. Used for short periods of restraint for non-painful procedures (e.g., blood collection) or before induction and gas anesthesia.

Drug	Dose	Route	Duration of Effect	Notes
Acepromazine	2 mg/kg	IM	30 minutes	Laryngeal reflexes are preserved.
Ketamine Xylazine	35 mg/kg 5 mg/kg	IM	30-60 minutes	First, inject acepromazine then xylazine, then ketamine into a different muscle.

- 5.5 Induction before inhalant anesthesia to facilitate intubation:

Drug	Dose	Route	Duration of Effect	Notes
Propofol	1-2 mg/kg	IV, slowly	Until discontinued	Administer to effect to facilitate intubation. It may cause respiratory depression.
Isoflurane	3% to 5%	Mask	Until discontinued	0.8 to 1.5 L/min. Initially, use a loose-fitting mask to minimize CO ₂ re-inhalation. Then switch to a tight-fitting mask.

- 5.6. Intubation:
 - 5.6.1. It is recommended to pre-oxygenate for 1 to 5 minutes with a tightly fitted mask with 100% oxygen to avoid apnea during induction and intubation.
 - 5.6.2. placement of an endotracheal tube is recommended for delivery of isoflurane anesthesia.
 - 5.6.3. Cuffed endotracheal tubes are preferred for animals weighing over 2kg as they reduce the possibility of aspiration of saliva or stomach contents.
 - 5.6.4. Intubation:
 - 5.6.4.1. Anesthetise with injectable anesthetics
 - 5.6.4.2. Lubricate endotracheal tube with sterile lubricant.

- 5.6.4.3. With the animal in sternal recumbency, extend the neck and head upward, so they are straight.
 - 5.6.4.4. Connect the tube to the stethoscope instead of the plate.
 - 5.6.4.5. Insert the tube into the mouth until it reaches resistance.
 - 5.6.4.6. Listen to the breath whistle (open of the larynx) and gently push the tube when the whistle sounds
 - 5.6.4.7. Confirm proper placement by checking for the animal's breath as it exits the endotracheal tube during exhalation.
 - 5.6.4.8. Secure the endotracheal tube by tying a piece of gauze around the tube and then behind the animal's head.
 - 5.6.4.9. Inflate the cuff of the endotracheal tube.
 - 5.6.4.10. Verify adequate ventilation of both lungs by auscultation.
- 5.7. Isoflurane anesthesia:
- 5.7.1. Induction:
 - 5.7.1.1. Adjust the oxygen flowmeter to 0.8 to 1.5 L/min.
 - 5.7.1.2. Adjust the isoflurane vaporizer to 3% to 5%.
 - 5.7.2. Maintenance:
 - 5.7.2.3. Adjust the flowmeter to 400 to 800mL/min.
 - 5.7.2.4. Adjust the isoflurane vaporizer to 1.5 to 3%.
 - 5.7.2.5. Monitor parameters (heart rate, oxygen saturation, respiration rate, temperature) every 5 minutes.
 - 5.7.3. Recovery:
 - 5.7.3.1. Turn off the isoflurane vaporizer but keep the animal on oxygen for 2 to 5 minutes or longer if oxygen saturation levels are low.
 - 5.7.3.2. Monitor temperature at the end of surgery and warm the animal up if necessary to speed up recovery.
 - 5.7.3.3. Deflate the endotracheal cuff and stay close to the animal if the animal is intubated.
 - 5.7.3.1. Remove the endotracheal tube as soon as the animal shows signs of impending arousal, i.e. when reflexes begin to return.
 - 5.7.3.2. Remove the IV catheter before placing the animal back in its cage.
 - 5.7.3.3. Monitor the animal's home cage to ensure it regains full consciousness.